Brain Research, 398 (1986) 425–428
Elsevier

Author Index

Abe, H., see Kita, T., 18
Alexander, G.M., see Schwartzman, R.J., 113
Anderson, W.W., Lewis, D.V., Swartzwelder, H.S. and Wilson, W.A., Magnesium-free medium activates seizure-like events in the rat hippocampal slice, 215
Applegate, C.D., Flashman, L.A. and Burchfiel, J.L., The effects of chronic desmethylimipramine on entorhinal cortical kindling in rats, 121
Archer, T., see Mohammed, A.K., 6
Arends, J.J.A. and Zeigler, H.P., Anatomical identification of an auditory pathway from a nucleus of the lateral lemniscal system to the frontal telencephalon (nucleus basalis) of the pigeon, 375
Argiolas, A., see Melis, M.R., 259

Barnes, C.L., see Itaya, S.K., 397
Bartlett, P.F., see Sinclair, C.M., 91
Bedran deCastro, J.C., see Khorram, O., 361
Benoist, J.M., Kayser, V., Gacel, G., Zajac, J.M., Gautron, M., Roques, B. and Guilbaud, G., Differential depressive action of two μ and δ opioid ligands on neuronal responses to noxious stimuli in the thalamic ventrobasal complex of rat, 49
Beresford, I.J.M., Clark, C.R. and Hughes, J., Measurement and characterization of neuronal cholecystokinin using a novel radioreceptor assay, 313
Berger, T.W., see Nisenbaum, E.S., 221
Berwald-Netter, Y., see Takeda, K., 75
Besson, J., Sarrieau, A., Vial, M., Marie, J.-C., Rosselin, G. and Rostène, W., Characterization and autoradiographic distribution of vasoactive intestinal peptide binding sites in the rat central nervous system, 329
Bloxham, C.A., see Whitford, C.A., 141
Botez, M.I., see Lalonde, R., 175
Bouldin, T.W., see Toews, A.D., 298
Boyd, Jr., F.T., Clarke, D.W. and Raizada, M.K., Insulin inhibits specific norepinephrine uptake in neuronal cultures from rat brain, 1
Briet, D., see Gourmelon, P., 128
Brusovanik, V.I., see Erin, A.N., 85
Buchwald, N.A., see Fisher, R.S., 148
Burchfiel, J.L., see Applegate, C.D., 121
Buzsáki, G., Hippocampal sharp waves: their origin and significance, 242

Candy, J.M., see Whitford, C.A., 141
Cardinali, D.P., see Rosenstein, R.E., 403
Castañeyra-Perdomo, A., see Meyer, G., 207
Cavalheiro, E.A., see Turski, L., 37
Chiba, T., see Kita, T., 18
Chihara, K., see Kita, T., 18
Christakos, C.M., see Windhorst, U., 181
Clark, C.R., see Beresford, I.J.M., 313
Clarke, D.W., see Boyd, Jr., F.T., 1
Colin, S. and Kruger, L., Peptidergic nociceptive axon visualization in whole-mount preparations of cornea and tympanic membrane in rat, 199
Coquerel, A., Dubuc, I., Menard, J.-F., Kitabgi, P. and Costentin, J., Naloxone-insensitive potentiation of neurotensin hypothermic effect by the enkephalinase inhibitor thiorphan, 386
Costentin, J., see Coquerel, A., 386
Court, L., see Gourmelon, P., 128
Crawley, J.M., Glowa, J.R., Majewska, M.D. and Paul, S.M., Anxiolytic activity of an endogenous adrenal steroid, 382
Critchlow, V., Dyke, A. and Kaler, L.W., Release of growth hormone, prolactin and somatostatin during perifusion of anterior pituitary and preoptic-medial basal hypothalamus from male and female rats, 347
Cruz, E.G., see Mobley, P.L., 366
Cuello, A.C., see Sofroniew, M.V., 393
Czuczwar, S.J., see Turski, L., 37

DeGroat, W.C., see Morgan, C., 11
De Paermentier, F., Heuschling, P., Knoops, B., Janssens de Varebeke, P. and Van den Bosch de Aguilar, P., A new model for quantification of microvascular regeneration after a lesion of the rat cerebral cortex, 419
De Toledo-Morrell, L., see Geinisman, Y., 266
Dubocovich, M.L., see Hensler, J.G., 407
Dubuc, I., see Coquerel, A., 386
Dyke, A., see Critchlow, V., 347

Ehinger, B.E., see Perez, M.T.R., 106
Eidelberg, E., see Schwartzman, R.J., 113
Enoka, R.M., see Windhorst, U., 181
Erin, A.M., Gorbunov, N.V., Brusovanik, V.I., Tyurin, V.A. and Prilipko, L.L., Stabilization of synaptic membranes by α-tocopherol against the damaging action of phospholipases. Possible mechanism of biological action of vitamin E, 85

Ferres-Torres, R., see Meyer, G., 207
Fisher, R.S., Buchwald, N.A., Hull, C.D. and Levine, M.S., The GABAergic striatonigral neurons of the cat: demonstration by double peroxidase labeling, 148
Flashman, L.A., see Applegate, C.D., 121
Fredholm, B.B., see Perez, M.T.R., 106
Freeman, J.A., see Langdon, R.B., 169
Fujita, T., see Kita, T., 18

Gacel, G., see Benoist, J.M., 49
Galindo-Mireles, D., see Meyer, G., 207
Gautron, M., see Benoist, J.M., 49
Geinisman, Y., De Toledo-Morrell, L. and Morrell, F., Aged rats need a preserved complement of perforated axospinous synapses per hippocampal neurons to maintain good spatial memory, 266
Gessa, G.L., see Melis, M.R., 259
Glowa, J.R., see Crawley, J.M., 382
Goines, N.D., see Toews, A.D., 298
Gonzalez-Hernandez, T., see Meyer, G., 207
Gorbunov, N.V., see Erin, A.M., 85
Gourmelon, P., Briet, D., Court, L. and Tsiang, H., Electrophysiological and sleep alterations in experimental mouse

rabies, 128
Govoni, S., see Lucchi, L., 212
Greig, D.I., see Sinclair, C.M., 91
Guilbaud, G., see Benoist, J.M., 49
Guitton, D., see Munoz, D.P., 185

Hamm, T.M., see Windhorst, U., 181
Hawkins, R.L. and Seeds, N.W., Effect of proteases and their inhibitors on neurite outgrowth from neonatal mouse sensory ganglia in culture, 63
Hensler, J.G. and Dubocovich, M.L., D_1-dopamine receptor activation mediates [^{3}H]acetylcholine release from rabbit retina, 407
Heuschling, P., see de Paermentier, F., 419
Hiramatsu, M., see Nabeshima, T., 354
Hirst, B.H., see Whitford, C.A., 141
Hochner, B. and Spira, M.E., Two distinct propagating regenerative potentials in a single ethanol-treated axon, 164
Hoffmann, G.E., see Watson, Jr., R.E., 157
Hong, J.S., see Li, S., 390
Hughes, J., see Beresford, I.J.M., 313
Hull, C.D., see Fisher, R.S., 148

Ikonomidou-Turski, C., see Turski, L., 37
Inagaki, S., see Yamano, M., 337
Itaya, S.K., Van Hoesen, G.W. and Barnes, C.L., Anterograde transsynaptic transport of WGA-HRP in the limbic system of rat and monkey, 397

Janssens de Varebeke, P., see De Paermentier, F., 419
Jeffrey, P.L., see Sinclair, C.M., 91
Jonsson, G., see Mohammed, A.K., 6

Kaji, H., see Kita, T., 18
Kaler, L.W., see Critchlow, V., 347
Kameyama, T., see Nabeshima, T., 354
Kato, M., see Matsuoka, S., 253
Katoh, A., see Nabeshima, T., 354
Kayser, V., see Benoist, J.M., 49
Khorram, O., Bedran deCastro, J.C. and McCann, S.M., The influence of suckling on the hypothalamic and pituitary secretion of immunoreactive α-melanocyte stimulating hormone, 361
Kita, T., Chihara, K., Abe, H., Minamitani, N., Kaji, H., Kodama, H., Chiba, T., Fujita, T. and Yanaihara, N., Regional distribution of gastrin-releasing peptide- and somatostatin-like immunoreactivity in the rabbit hypothalamus, 18
Kitabgi, P., see Coquerel, A., 386
Kitamura, K., see Matsuoka, S., 253
Kito, S., see Yamano, M., 337
Knoops, B., see De Paermentier, F., 419
Kodama, H., see Kita, T., 18
Koehler, W., see Windhorst, U., 181
Koenig, E., Isolation of native Mauthner cell axoplasm and an analysis of organelle movement in non-aqueous and aqueous media, 288
Kruger, L., see Colin, S., 199

Lalonde, R., Delayed spontaneous alternation in weaver mutant mice, 178
Lalonde, R. and Botez, M.I., Navigational deficits in weaver mutant mice, 175
Langdon, R.B. and Freeman, J.A., Antagonists of glutaminergic neurotransmission block retinotectal transmission in goldfish, 169
Levine, M.S., see Fisher, R.S., 148
Lewis, D.V., see Anderson, W.W., 215
Li, S., Sivam, S.P. and Hong, J.S., Regulation of the concentration of dynorphin A_{1-8} in the striatonigral pathway by the dopaminergic system, 390
Lind, A., see van Marle, J., 191
Lindström, K., see Perez, M.T.R., 106
Lucchi, L., Govoni, S., Spano, P.F. and Trabucchi, M., Release of Met-enkephalin from rat striatal slices: effect of amphetamine and fipexide, 212

Majewska, M.D., see Crawley, J.N., 382
Marie, J.-C., see Besson, J., 329
Matsuoka, S., Shima, F., Suzuki, M., Kitamura, K. and Kato, M., Which is responsible for circling behavior in rats, locus coeruleus or dorsolateral pontine tegmentum? 253
Matsuzaki, T., see Yamano, M., 337
McCann, S.M., see Khorram, O., 361
McCaslin, P.P. and Morgan, W.W., 2-Amino-7-phosphonoheptanoic acid, a selective N-methyl-D-aspartate antagonist, blocks swim-induced elevation of cerebellar cyclic guanosine monophosphate, 71
McEwen, B.S., see Meaney, M.J., 324
McNamara, J.O., see Shin, C., 370
Meaney, M.J. and McEwen, B.S., Testosterone implants into the amygdala during the neonatal period masculinize the social play of juvenile female rats, 324
Meldrum, B.S., see Millan, M.H., 204
Melis, M.T., Argiolas, A. and Gessa, G.L., Oxytocin-induced penile erection and yawning: site of action in the brain, 259
Menard, J.-F., see Coquerel, A., 386
Meyer, G., Galindo-Mireles, D., Gonzalez-Hernandez, T., Castañeyra-Perdomo, A. and Ferres-Torres, R., Direct projections from the reticular formation of the medulla oblongata to the anterior cingulate cortex in the mouse and the rat, 207
Millan, M.H., Patel, S. and Meldrum, B.S., Olfactory bulbectomy protects against pilocarpine-induced motor limbic seizures in rats, 204
Minamitani, N., see Kita, T., 18
Mizuno, N., see Sugimoto, T., 195
Mobley, P.L., Scott, S.L. and Cruz, E.G., Protein kinase C in astrocytes: a determinant of cell morphology, 366
Mohammed, A.K., Jonsson, G. and Archer, T., Selective lesioning of forebrain noradrenaline neurons at birth abolishes the improved maze learning performance induced by rearing in complex environment, 6
Moran, J., Pasantes-Morales, H. and Redburn, D.A., Glutamate receptor agonists release [^{3}H]GABA preferentially from horizontal cells, 276
Morgan, C., Nadelhaft, I. and deGroat, W.C., The distribution within the spinal cord of visceral primary afferent axons carried by the lumbar colonic nerve of the cat, 11
Morgan, W.W., see McCaslin, P.P., 71
Morrell, F., see Geinisman, Y., 266
Munoz, D.P. and Guitton, D., Presaccadic burst discharges of tecto-reticulo-spinal neurons in the alert head-free and -fixed cat, 185

Nabeshima, T., Katoh, A., Hiramatsu, M. and Kameyama, T., A role played by dopamine and opioid neuronal systems in stress-induced motor suppression (conditioned suppression of motility) in mice, 354
Nadelhaft, I., see Morgan, C., 11

Nisenbaum, E.S., Stricker, E.M., Zigmond, M.J. and Berger, T.W., Long-term effects of dopamine-depleting brain lesions on spontaneous activity of Type II striatal neurons: relation to behavioral recovery, 221

Oishi, R., see Yoshitomi, I., 57

Pasantes-Morales, H., see Moran, J., 276
Patel, S., see Millan, M.H., 204
Paul, S.M., see Crawley, J.N., 382
Pearson, R.C.A., see Sofroniew, M.V., 393
Perez, M.T.R., Ehinger, B.E., Lindström, K. and Fredholm, B.B., Release of endogenous and radioactive purines from the rabbit retina, 106
Prilipko, L.L., see Erin, A.N., 85

Raizada, M.K., see Boyd, Jr., F.T., 1
Ray, R.B., see Toews, A.D., 298
Redburn, D.A., see Moran, J., 276
Rigsbee, L.C., see Shin, C., 370
Roques, B., see Benoist, J.M., 49
Rosenstein, R.E. and Cardinali, D.P., Melatonin increases in vivo GABA accumulation in rat hypothalamus, cerebellum, cerebral cortex and pineal gland, 403
Rosselin, G., see Besson, J., 329
Rostène, W., see Besson, J., 329

Saeki, K., see Yoshitomi, I., 57
Sarrieau, A., see Besson, J., 329
Schmechel, D.E., see Walsh, T.J., 23
Schneiderman, J.H., Low concentrations of penicillin reveal rhythmic, synchronous synaptic potentials in hippocampal slice, 231
Schulz, D.W., see Walsh, T.J., 23
Schwartzman, R.J., Eidelberg, E. and Alexander, G.M., Asymmetrical regional changes in energy metabolism of the central nervous system during walking, 113
Scott, S.L., see Mobley, P.L., 366
Seeds, N.W., see Hawkins, R.L., 63
Shima, F., see Matsuoka, S., 253
Shimahara, T., see Takeda, K., 75
Shin, C., Rigsbee, L.C. and McNamara, J.O., Anti-seizure and anti-epileptogenic effect of γ-vinyl γ-aminobutyric acid in amygdaloid kindling, 370
Shinohara, Y., see Yamano, M., 337
Sieklucka-Dziuba, M., see Turski, L., 37
Sinclair, C.M., Bartlett, P.F., Greig, D.I. and Jeffrey, P.L., Distribution of Thy-1 in the avain nervous system: immunohistochemical and absorption analyses with a monoclonal antibody, 91
Sivam, S.P., see Li, S., 390
Snell, C.R., see Whitford, C.A., 141
Sofroniew, M.V., Pearson, R.C.A., Cuello, A.C., Tagari, P.C. and Stephens, P.H., Parenterally administered GM_1 ganglioside prevents retrograde degeneration of cholinergic cells of the rat basal forebrain, 393
Spano, P.F., see Lucchi, L., 212
Spira, M.E., see Hochner, B., 164
Stephens, P.H., see Sofroniew, M.V., 393
Stricker, E.M., see Nisenbaum, E.S., 221
Stuart, D.G., see Windhorst, U., 181
Sugimoto, T. and Mizuno, N., Immunohistochemical demonstration of neurotensin in striatal neurons of the cat, with particular reference to coexistence with enkephalin, 195
Suzuki, M., see Matsuoka, S., 253
Suzuki, Y. and Taguchi, K., In vivo voltammetric studies of the effect of morphine on the serotonergic system in the cat spinal cord, 413
Swartzwelder, H.S., see Anderson, W.W., 215

Tagari, P.C., see Sofroniew, M.V., 393
Taguchi, K., see Suzuki, Y., 413
Takeda, K., Shimahara, T., Tauc, L. and Berwald-Netter, Y., Acetylcholine-activated currents in mouse neuroblastoma cells, 75
Tauc, L., see Takeda, K., 75
Tilson, H.A., see Walsh, T.J., 23
Toews, A.D., Ray, R.B., Goines, N.D. and Bouldin, T.W., Increased synthesis of membrane macromolecules is an early response of retinal neurons to trimethyltin intoxication, 298
Tohyama, M., see Yamano, M., 337
Trabucchi, M., see Lucchi, L., 212
Tsiang, H., see Gourmelon, P., 128
Turski, L., Cavalheiro, E.A., Sieklucka-Dziuba, M., Ikonomidou-Turski, C., Czuczwar, S.J. and Turski, W.A., Seizures produced by pilocarpine: neuropathological sequelae and activity of glutamate decarboxylase in the rat forebrain, 37
Turski, W.A., see Turski, L., 37
Tyurin, V.A., see Erin, A.N., 85

Van den Bosch de Aguilar, P., see de Paermentier, F., 419
Van Hoesen, G.W., see Itaya, S.K., 397
Van Marle, J., Veldsema-Currie, R.D., Lind, A. and van Weeren-Kramer, J., Dexamethasone reduces the high-affinity binding of hemicholinium-3 to rat phrenic nerve endings in vitro: an autoradiographic study, 191
Van Weeren-Kramer, J., see van Marle, J., 191
Veldsema-Currie, R.D., see van Marle, J., 191
Vial, M., see Besson, J., 329
Walsh, T.J., Schulz, D.W., Tilson, H.A. and Schmechel, D.E., Colchicine-induced granule cells loss in rat hippocampus: selective behavioral and histological alterations, 23

Watson, Jr., R.E., Hoffman, G.E. and Wiegand, S.J., Sexually dimorphic opioid distribution in the preoptic area: manipulation by gonadal steroids, 157
Weiland, N.G. and Wise, P.M., Effects of age on β_1- and β_2-adrenergic receptors in the brain assessed by quantitative autoradiography, 305
Whitford, C.A., Bloxham, C.A., Snell, C.R., Candy, J.M. and Hirst, B.H., Regional distribution of high-affinity [^{3}H]somatostatin binding sites in the human brain, 141
Wiegand, S.J., see Watson, Jr., R.E., 157
Wilson, W.A., see Anderson, W.W., 215
Windhorst, U., Christakos, C.N., Koehler, W., Hamm, T.M., Enoka, R.M. and Stuart, D.G., Amplitude reduction of motor unit twitches during repetitive activation is accompanied by relative increase of hyperpolarizing membrane potential trajectories in homonymous α-motoneurons, 181
Wise, P.M., see Weiland, N.G., 305

Yamano, M., Inagaki, S., Kito, S., Matsuzaki, T., Shinohara, Y. and Tohyama, M., Enkephalinergic projection from the ventromedial hypothalamic nucleus to the midbrain central gray matter in the rat: an immunocytochemical analysis, 337
Yanaihara, N., see Kita, T., 18
Yoshitomi, I., Oishi, R. and Saeki, K., Involvement of opioid and non-opioid mechanisms in footshock-induced enhance-

ment of brain histamine turnover in mice, 57

Zajac, J.M., see Benoist, J.M., 49

Zeigler, H.P., see Arends, J.J.A., 375
Zigmond, M.J., see Nisenbaum, E.S., 221

(contents continued)

Anti-seizure and anti-epileptogenic effect of γ-vinyl γ-aminobutyric acid in amygdaloid kindling (BRE 21847)
by C. Shin, L.C. Rigsbee and J.O. McNamara (U.S.A.) 370
Anatomical identification of an auditory pathway from a nucleus of the lateral lemniscal system to the frontal telencephalon (nucleus basalis) of the pigeon (BRE 21857)
by J.J.A. Arends and H.P. Zeigler (U.S.A.) 375
Anxiolytic activity of an endogenous adrenal steroid (BRE 21888)
by J.N. Crawley, J.R. Glowa, M.D. Majewska and S.M. Paul (U.S.A.) 382
Naloxone-insensitive potentiation of neurotensin hypothermic effect by the enkephalinase inhibitor thiorphan (BRE 21890)
by A. Coquerel, I. Dubuc, J.-F. Menard, P. Kitabgi and J. Costentin (France) 386
Regulation of the concentration of dynorphin A_{1-8} in the striatonigral pathway by the dopaminergic system (BRE 21894)
by S. Li, S.P. Sivam and J.S. Hong (U.S.A.) 390
Parenterally administered GM_1 ganglioside prevents retrograde degeneration of cholinergic cells of the rat basal forebrain (BRE 21896)
by M.V. Sofroniew, R.C.A. Pearson, A.C. Cuello, P.C. Tagari and P.H. Stephens (U.K. and Canada) ... 393
Anterograde transsynaptic transport of WGA-HRP in the limbic system of rat and monkey (BRE 21897)
by S.K. Itaya, G.W. Van Hoesen and C.L. Barnes (U.S.A.) 397
Melatonin increases in vivo GABA accumulation in rat hypothalamus, cerebellum, cerebral cortex and pineal gland (BRE 21900)
by R.E. Rosenstein and D.P. Cardinali (Argentina) 403
D_1-dopamine receptor activation mediates [^{3}H]acetylcholine release from rabbit retina (BRE 21902)
by J.G. Hensler and M.L. Dubocovich (U.S.A.) 407
In vivo voltammetric studies of the effect of morphine on the serotonergic system in the cat spinal cord (BRE 21904)
by Y. Suzuki and K. Taguchi (Japan) 413
A new model for quantification of microvascular regeneration after a lesion of the rat cerebral cortex (BRE 21905)
by F. de Paermentier, P. Heuschling, B. Knoops, P. Janssens de Varebeke and P. van den Bosch de Aguilar (Belgium) 419

Author Index 425